arXiv:1511.01847v2 [math.AG] 26 May 2016

ON THE SINGULAR SHEAVES IN THE FINE SIMPSON MODULI SPACES OF 1-DIMENSIONAL SHEAVES

OLEKSANDR IENA AND ALAIN LEYTEM

Abstract. In the Simpson moduli space M of semi-stable sheaves with Hilbert polynomial $dm-1$ on a projective plane we study the closed subvariety M' of sheaves that are not locally free on their support. We show that for $d \geqslant 4$ it is a singular subvariety of codimension 2 in M. The blow up of M along M' is interpreted as a (partial) modification of $M \setminus M'$ by vector bundles (on support).

Contents

1. Introduction 1
2. Basic constructions 2
3. Singular sheaves 4
4. Modifying the boundary by vector bundles 7
Appendix A. On the ideals of points on planar curves 10
References 11

1. Introduction

Let $\Bbbk$ be an algebraically closed field of characteristic zero, let V be a vector space over $\Bbbk$ of dimension 3, and let $\mathbb{P}_2 = \mathbb{P}V$ be the corresponding projective plane. Consider a linear polynomial with integer coefficients $P(m) = dm + c \in \mathbb{Z}[m]$, $d \in \mathbb{Z}_{>0}$, $c \in \mathbb{Z}$. Let $M := M_P(\mathbb{P}_2)$ be the Simpson coarse moduli space (cf. [9]) of semi-stable sheaves on $\mathbb{P}_2$ with Hilbert polynomial P. As shown in [6], M is an irreducible locally factorial variety of dimension d^2+1, smooth if $\gcd(d,c) = 1$.

Singular sheaves. The sheaves from M are torsion sheaves on $\mathbb{P}_2$, they are supported on curves of degree d. Restricted to their (Fitting) support most of sheaves in M are vector bundles on curves. Such sheaves constitute an open subvariety M_B of M. Its complement M', the closed subvariety of sheaves that are not locally free on their support, is in general non-empty. This way one could consider M as a compactification of M_B. We call the sheaves from the boundary $M' = M \setminus M_B$ *singular*.

The boundary M' does not have the minimal codimension in general. Loosely speaking, one glues together too many different directions at infinity. For example, for $c \in \mathbb{Z}$ with $\gcd(3,c) = 1$, all moduli spaces M_{3m+c} are isomorphic to the universal plane cubic curve and M'_{3m+c} is a smooth subvariety of codimension 2 isomorphic to the universal singular locus of a cubic curve (cf. [5]).

The main result. As demonstrated in [4], for $P(m) = 4m+c$ with $\gcd(4,c) = 1$, the subvariety M' is singular of codimension 2. The main result of this paper is the following generalization of [4].

2010 *Mathematics Subject Classification.* 14D20.
Key words and phrases. Simpson moduli spaces, coherent sheaves, vector bundles on curves, singular sheaves.

Theorem 1.1. *For an integer $d \geqslant 4$, let $M = M_{dm-1}(\mathbb{P}_2)$ be the Simpson moduli space of (semi-)stable sheaves on $\mathbb{P}_2$ with Hilbert polynomial $dm-1$. Let $M' \subseteq M$ be the subvariety of singular sheaves. Then M' is singular of codimension 2.*

Using our understanding of M' and the construction from [5] we obtain as a consequence Theorem 4.7, which allows to interpret the blow up of M along M' as a modification of the boundary M' by a divisor consisting generically of vector bundles.

Structure of the paper. In Section 2 we recall the results from [8] and identify the open subvariety M_0 in M of sheaves without global sections with an open subvariety of a projective bundle $\mathbb{B}$ over a variety of Kronecker modules N. In Section 3, using a convenient characterization of free ideals of fat curvilinear points on planar curves from Appendix A, we show that a generic fibre of $M_0' = M_0 \cap M'$ over N is a union of projective subspaces of codimension 2 in the fibre of $\mathbb{B}$, thus it is singular of codimension two, which allows to prove the main result. In Section 4 we briefly discuss how our analysis can be used for modifying the boundary M' by vector bundles (on support).

2. Basic constructions

2.1. **Kronecker modules.** Let $\mathbb{V}$ be the affine space of Kronecker modules

$$(n-1)\mathcal{O}_{\mathbb{P}_2}(-1) \xrightarrow{\Phi} n\mathcal{O}_{\mathbb{P}_2}. \tag{1}$$

There is a natural group action of $G = (\mathrm{GL}_{n-1}(\Bbbk) \times \mathrm{GL}_n(\Bbbk))/\Bbbk^*$ on $\mathbb{V}$. Since $\gcd(n-1, n) = 1$, all semistable points of this action are stable and G acts freely on the open on the open subset of stable points $\mathbb{V}^s$. Then $\Phi \in \mathbb{V}^s$ if and only if Φ does not lie in the same orbit with a Kronecker module with a zero block of size $j \times (n-j)$, $1 \leqslant j \leqslant n-1$ (cf. [1, Proposition 15], [3, 6.2]). In particular, Kronecker modules with linear independent maximal minors are stable. There exists a geometric quotient $N = N(3; n-1, n) = \mathbb{V}^s/G$, which is a smooth projective variety of dimension $(n-1)n$. For more details consult [3, Section 6] and [1, Section III].

The cokernel of a stable Kronecker module $\Phi \in \mathbb{V}^s$ is an ideal of a zero-dimensional scheme Z of length $(n-1)n/2$ if the maximal minors of Φ are coprime. In this case the maximal minors $d_0, \ldots, d_{n-1}$ are linear independent and there is a resolution

$$0 \to (n-1)\mathcal{O}_{\mathbb{P}_2}(-n) \xrightarrow{\Phi} n\mathcal{O}_{\mathbb{P}_2}(-n+1) \xrightarrow{\begin{pmatrix} d_0 \\ \vdots \\ d_{n-1} \end{pmatrix}} \mathcal{O}_{\mathbb{P}_2} \to \mathcal{O}_Z \to 0. \tag{2}$$

Moreover Z does not lie on a curve of degree $n-2$. Let $\mathbb{V}_0$ denote the open subvariety of $\Phi \in \mathbb{V}^s$ of Kronecker modules with coprime maximal minors. Let $N_0 \subseteq N$ be the corresponding open subvariety in the quotient space.

This way one obtains a morphism from $N_0 \subseteq N$ to the Hilbert scheme of zero-dimensional subschemes of $\mathbb{P}_2$ of length $l = (n-1)n/2$, which sends a class of $\Phi \in \mathbb{V}^s$ to the zero scheme of its maximal minors. Since every zero dimensional scheme of length l that does not lie on a curve of degree $n-2$ has a minimal resolution of type (2), this gives an isomorphism of N_0 and the open subvariety $H_0 \subseteq H = \mathbb{P}_2^{[l]}$ consisting of Z that do not lie on a curve of degree $n-2$. The complement of H_0 is an irreducible hypersurface (as mentioned e. g. in [2, p. 46]).

2.2. **Projective bundle over N.** Let $\mathbb{U}_1 = (n-1)\Gamma(\mathbb{P}_2, \mathcal{O}_{\mathbb{P}_2}(1))$, $\mathbb{U}_2 = n\Gamma(\mathbb{P}_2, \mathcal{O}_{\mathbb{P}_2}(2))$. Consider the trivial vector bundles $\mathbb{V} \times \mathbb{U}_1$ and $\mathbb{V} \times \mathbb{U}_2$ over $\mathbb{V}$. Consider the following morphism between them:

$$\mathbb{V} \times \mathbb{U}_1 \xrightarrow{F} \mathbb{V} \times \mathbb{U}_2, \quad (\Phi, L) \mapsto (\Phi, L \cdot \Phi).$$

Lemma 2.1. *The morphism F is injective over $\mathbb{V}^s$.*

Proof. Let $\Phi \in \mathbb{V}^s$ and assume $L \cdot \Phi = 0$ for some non-zero $0 \neq L = (l_1, \dots, l_{n-1}) \in \mathbb{U}_1$.

Since $l_i \in \Gamma(\mathbb{P}_2, \mathcal{O}_{\mathbb{P}_2}(1))$, the dimension of the vector space generated by $\{l_i\}_i$ is at most 3. So for some $B \in \mathrm{GL}_{n-1}(\Bbbk)$ we get $L \cdot B = L' = (l'_1, l'_2, l'_3, 0, \dots, 0)$ such that the first non-zero entries are linear independent. Then, since Φ is semistable if and only if $B^{-1}\Phi$ is semistable and since $L\Phi = (LB) \cdot (B^{-1}\Phi) = 0$, we may assume without loss of generality that $L = (l_1, l_2, l_3, 0, \dots, 0)$ and the first non-zero entries of L are linear independent.

If $l_1 \neq 0$, $l_2 = l_3 = 0$, then $L\Phi = 0$ implies that the first row of Φ is zero, which contradicts the stability of Φ.

If $l_1 \neq 0$, $l_2 \neq 0$, $l_3 = 0$, then the syzygy module of (l_1, l_2) is generated by $\left(\begin{smallmatrix} l_2 \\ -l_1 \end{smallmatrix}\right)$. So the columns of the first two rows of Φ are scalar multiples of $\left(\begin{smallmatrix} l_2 \\ -l_1 \end{smallmatrix}\right)$ and hence, after performing elementary transformations on the columns of Φ, Φ is equivalent to a matrix with a zero block of size $2 \times (n-1)$, which contradicts the stability of Φ.

If $l_i \neq 0$, $i = 1, 2, 3$, then the syzygy module of (l_1, l_2, l_3) is generated by 3 linear independent generators

$$\begin{pmatrix} 0 \\ l_3 \\ -l_2 \end{pmatrix}, \begin{pmatrix} -l_3 \\ 0 \\ l_1 \end{pmatrix}, \begin{pmatrix} l_2 \\ -l_1 \\ 0 \end{pmatrix},$$

which implies that Φ is equivalent to a matrix with a zero block of size $3 \times (n-3)$, which contradicts the stability of Φ. □

Therefore, $\mathbb{V}^s \times \mathbb{U}_1 \xrightarrow{F} \mathbb{V}^s \times \mathbb{U}_2$ is a vector subbundle and hence the cokernel of F is a vector bundle of rank $6 \cdot n - 3 \cdot (n-1) = 3n + 3 = 3d$ denoted by E.

The group action of $\mathrm{GL}_{n-1}(\Bbbk) \times \mathrm{GL}_n(\Bbbk)$ on $\mathbb{V}^s \times \mathbb{U}_1$ and $\mathbb{V}^s \times \mathbb{U}_2$ induces a group action of $\mathrm{GL}_{n-1}(\Bbbk) \times \mathrm{GL}_n(\Bbbk)$ on E and hence an action of $G = (\mathrm{GL}_{n-1}(\Bbbk) \times \mathrm{GL}_n(\Bbbk))/\Bbbk^*$ on $\mathbb{P}E$. Finally, since the stabilizer of $\Phi \in \mathbb{V}^s$ under the action of G is trivial, it acts trivially on the fibres of $\mathbb{P}E$ and thus $\mathbb{P}E$ descends to a projective $\mathbb{P}_{3n+2}$-bundle

$$\mathbb{B} \xrightarrow{\nu} N = N(3; n-1, n) = \mathbb{V}^s/G.$$

Let $\mathbb{W}$ be the affine variety of morphisms

$$\mathcal{O}_{\mathbb{P}_2}(-3) \oplus (n-1)\mathcal{O}_{\mathbb{P}_2}(-2) \xrightarrow{A} n\mathcal{O}_{\mathbb{P}_2}(-1). \tag{3}$$

Notice that $\mathbb{W}$ can be identified with $\mathbb{V} \times \mathbb{U}_2$ by the isomorphism

$$\mathbb{V} \times \mathbb{U}_2 \to \mathbb{W}, \quad (\Phi, Q) \mapsto \begin{pmatrix} Q \\ \Phi \end{pmatrix}.$$

The group

$$G' = (\mathrm{Aut}(\mathcal{O}_{\mathbb{P}_2}(-3) \oplus (n-1)\mathcal{O}_{\mathbb{P}_2}(-2)) \times \mathrm{Aut}(n\mathcal{O}_{\mathbb{P}_2}(-1))/\Bbbk^*.$$

acts on $\mathbb{W} \cong \mathbb{V} \times \mathbb{U}_2$. As shown in [8, Proposition 7.7], $\mathbb{B}$ is a geometric quotient of $\mathbb{V}^s \times \mathbb{U}_2 \setminus \mathrm{Im}\, F$ with respect to G'.

2.3. **Moduli space** $M_{dm-1}(\mathbb{P}_2)$**.** Let $d \geqslant 4$, $d = n + 1$, be an integer. Let $M = M_{dm-1}(\mathbb{P}_2)$ be the Simpson moduli space of (semi-)stable sheaves on $\mathbb{P}_2$ with Hilbert polynomial $dm - 1$. In [8] is was shown that M contains an open dense subvariety M_0 of isomorphism classes of sheaves $\mathcal{F}$ with $h^0(\mathcal{F}) = 0$.

2.3.1. *Sheaves in* M_0. By [8, Claim 4.2] the sheaves in M without global sections are exactly the cokernels of the injective morphisms (3) with $A = \left(\begin{smallmatrix} Q \\ \Phi \end{smallmatrix}\right)$, $\det A \neq 0$, $\Phi \in \mathbb{V}^s$. This allows to describe M_0 as an open subvariety in $\mathbb{B}$.

Let $\mathbb{W}_0$ be the open subvariety of injective morphisms in $\mathbb{W}$ parameterizing the points from M_0. Since the determinant of a matrix from the image of F is zero, one sees that

$$\mathbb{W}_0 \subseteq \mathbb{V}^s \times \mathbb{U}_2 \setminus \mathrm{Im}\, F,$$

which allows to conclude that $M_0 = \mathbb{W}_0/G'$ is an open subvariety of $\mathbb{B}$.

Let $\mathbb{B}_0 = \mathbb{B}|_{N_0}$ be the restriction of $\mathbb{B}$ to the open subscheme $N_0 \subseteq N$.

From the exact sequence (2) it follows that a matrix $A = \binom{Q}{\Phi} \in \mathbb{W}$, $\Phi \in \mathbb{V}_0$, has zero determinant if and only if A lies in the image of F. Therefore, the fibres of $\mathbb{B}$ over N_0 are contained in M_0 and thus $\mathbb{B}_0 \subseteq M_0$. As shown in [10] the codimension of the complement of $\mathbb{B}_0$ in M is at least 2.

2.3.2. *Sheaves in* $\mathbb{B}_0$. Sheaves in $\mathbb{B}_0$ are exactly the twisted ideal sheaves $\mathcal{I}_{Z\subseteq C}(d-3)$ of zero dimensional schemes Z of length l lying on a curve C of degree d such that Z is not contained in a curve of degree $d-3$. In other words the sheaves $\mathcal{F}$ in $\mathbb{B}_0$ are given by the short exact sequences

$$0 \to \mathcal{F} \to \mathcal{O}_C(d-3) \to \mathcal{O}_Z \to 0$$

with $Z \subseteq C$ as described.

A fibre over a point $[\Phi]$ from N_0 can be seen as the space of plane curves of degree d through the corresponding subscheme of l points. The identification is given by the map

$$\nu^{-1}([\Phi]) \ni [A] \mapsto \langle \det A \rangle \in \mathbb{P}(S^d V^*).$$

Indeed, if two matrices over $[\Phi]$ are equivalent, then their determinants are equal up to a non-zero constant multiple and hence the map above is well-defined. On the other hand, if two matrices $A = \binom{Q}{\Phi}$ and $A' = \binom{Q'}{\Phi}$ have equal determinants, then $Q - Q'$ lies in the syzygy module of the maximal minors of Φ and hence (cf. (2)) is a linear combination of the rows of Φ, which means that A and A' are equivalent.

3. Singular sheaves

Let M' be the subvariety of singular sheaves in M and let $M'_0 = M' \cap M_0$. Let us consider the restriction of ν to M'_0 and describe some of its fibres.

3.1. **Generic fibres.** Let N_c be the open subset of N_0 that corresponds to l different points. Under the isomorphism $N_0 \cong H_0$ it corresponds to the open subvariety $H_c \subseteq H_0$ of the configurations of l points on $\mathbb{P}_2$ that do not lie on a curve of degree $d-3$.

Let $[\Phi] \in N_c$ and let $Z = \{pt_1, \dots, pt_l\}$ be the corresponding zero-dimensional scheme.

Let $m_1, \dots, m_l$ be the monomials of degree $d-3$ in variables x_0, x_1, x_2 ordered, say, in lexicographical order:

$$m_1 = x_0^{d-3}, \quad m_2 = x_0^{d-4}x_1, \quad m_3 = x_0^{d-4}x_2, \quad \dots$$

Since Z does not lie on a curve of degree $d-3$, the matrix

$$\begin{pmatrix} m_1(pt_1) & \dots & m_l(pt_1) \\ \vdots & \ddots & \vdots \\ m_1(pt_l) & \dots & m_l(pt_l) \end{pmatrix}$$

has full rank. Assume without loss of generality that $pt_1 = \langle 1, 0, 0 \rangle$. Then the matrix is

$$\begin{pmatrix} 1 & 0 & \dots & 0 \\ m_1(pt_2) & m_2(pt_2) & \dots & m_l(pt_2) \\ \vdots & \vdots & \ddots & \vdots \\ m_1(pt_l) & m_2(pt_l) & \dots & m_l(pt_l) \end{pmatrix}$$

and therefore the matrix

$$\begin{pmatrix} m_1(pt_2) & m_2(pt_2) & \dots & m_l(pt_2) \\ \vdots & \vdots & \ddots & \vdots \\ m_1(pt_l) & m_2(pt_l) & \dots & m_l(pt_l) \end{pmatrix}$$

has full rank and thus there exists a homogeneous polynomial q of degree $d-3$ vanishing at the points $pt_2, \dots, pt_l$ with the coefficient 1 in front of the monomial $m_1 = x_0^{d-3}$. Therefore, the forms $x_0^2x_1q$ and $x_0^2x_2q$ vanish at Z. Notice that $x_0^2x_1q$ has the monomial $x_0^{d-1}x_1$ but does not have $x_0^{d-1}x_2$, and $x_0^2x_2q$ has $x_0^{d-1}x_2$ but does not have $x_0^{d-1}x_1$.

Let $\mathcal{E}$ be a sheaf over $[\Phi]$. By Lemma A.1 it is singular at pt_1 if and only if pt_1 is a singular point of the support C of $\mathcal{E}$. The latter holds if and only if in the homogeneous polynomial of degree d defining C the coefficients in front of the monomials $x_0^{d-1}x_1$ and $x_0^{d-1}x_2$ vanish. Therefore, taking into account the considerations above, sheaves over $[\Phi]$ singular at pt_1 constitute a projective subspace of codimension 2 in the fibre $F := \nu^{-1}([\Phi]) \cong \mathbb{P}_{3d-1}$. Since our argument can be repeated for each point pt_i, $i = 1, \dots, l$, we conclude that the sheaves over $[\Phi]$ singular at pt_i constitute a projective subspace F_i of codimension 2 in the fibre $\nu^{-1}([\Phi]) \cong \mathbb{P}_{3d-1}$.

Now let us clarify how the linear subspaces in the fibre corresponding to different points pt_i intersect with each other. First of all notice that Z must contain a triple of non-collinear points because otherwise Z must lie on a line. Therefore, in addition to the assumption $pt_1 = \langle 1, 0, 0\rangle$, we can assume without loss of generality that $pt_2 = \langle 0, 1, 0\rangle$, $pt_3 = \langle 0, 0, 1\rangle$. Then the conditions for being singular at these three different points read as the absence of the following monomials in the equation of C:

$$\begin{aligned} &x_0^{d-1}x_1, x_0^{d-1}x_2 && \text{for the point } pt_1;\\ &x_1^{d-1}x_0, x_1^{d-1}x_2 && \text{for the point } pt_2;\\ &x_2^{d-1}x_0, x_2^{d-1}x_1 && \text{for the point } pt_3. \end{aligned} \tag{4}$$

These conditions are clearly independent of each other. Moreover, these conditions are independent from the conditions imposed on the curves of degree d by the requirement $Z \subseteq C$. This is the case since the forms $x_0^2x_1q$ and $x_0^2x_2q$ constructed above clearly do not have monomials $x_1^{d-1}x_0$, $x_1^{d-1}x_2$, $x_2^{d-1}x_0$, $x_2^{d-1}x_1$. This produces a way to construct the curves through Z with only one of the monomials from (4).

We conclude that for every pair of different indices $\operatorname{codim}_F F_i \cap F_j = 4$. Moreover, for every triple of different indices i, j, ν corresponding to 3 non-collinear points pt_i, pt_j, pt_ν from Z $\operatorname{codim}_F F_i \cap F_j \cap F_\nu = 6$. Finally we obtain the following.

Lemma 3.1. *The fibres of M_0' over N_c are unions of l different linear subspaces of $\nu^{-1}([\Phi]) \cong \mathbb{P}_{3d-1}$ of codimension 2 such that each pair intersects in codimension 4 and each triple corresponding to 3 non-collinear points intersects in codimension 6. In particular the fibres are singular.*

Remark 3.2. One can show that $\operatorname{codim}_F F_i \cap F_j \cap F_\nu = 6$ for every triple of different indices i, j, ν corresponding to 3 different points pt_i, pt_j, pt_ν from Z.

Remark 3.3. In general it is not true that F_i intersect transversally. For example, for $d = 6$ and $Z = \{pt_1, \dots, pt_{10}\}$ with

$$\begin{aligned} &pt_1 = \langle 1, 0, 0\rangle, && pt_2 = \langle 0, 1, 0\rangle, && pt_3 = \langle 0, 0, 1\rangle, && pt_4 = \langle 0, 1, 1\rangle, && pt_5 = \langle 0, 1, -1\rangle,\\ &pt_6 = \langle 1, -2, 0\rangle, && pt_7 = \langle 1, 2, -1\rangle, && pt_8 = \langle 1, 1, -2\rangle, && pt_9 = \langle 1, -1, 1\rangle, && pt_{10} = \langle 1, 1, -1\rangle, \end{aligned}$$

$\operatorname{codim}_F F_1 \cap F_2 \cap F_3 \cap F_4 = 8$ but $\operatorname{codim}_F F_1 \cap F_2 \cap F_3 \cap F_4 \cap F_5 = 9$.

3.2. **Fibres over N_1.** Let N_1 be the open subset of $N_0 \setminus N_c$ that corresponds to $l-2$ different simple points and one double point.

Let $[\Phi] \in N_1$ and let $Z = \{pt_1\} \cup \{pt_2, \dots, pt_{l-1}\}$ be the corresponding zero-dimensional scheme, where pt_1 is a double point. Without loss of generality, applying if necessary a coordinate change, we may assume that pt_1 is given by the ideal (x_1^2, x_2).

3.2.1. *Sheaves singular at the double point.* Since Z does not lie on a curve of degree $d-3$, the matrix

$$\begin{pmatrix} 1 & 0 & 0 & \dots & 0 \\ 0 & 1 & 0 & \dots & 0 \\ m_1(pt_2) & m_2(pt_2) & m_3(pt_2) & \dots & m_l(pt_2) \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ m_1(pt_{l-1}) & m_2(pt_{l-1}) & m_3(pt_{l-1}) & \dots & m_l(pt_{l-1}) \end{pmatrix}$$

has full rank. Therefore, there exist homogeneous polynomials q and q' of degree $d-3$ vanishing at the points $pt_2, \dots, pt_{l-1}$ such that q does not have monomial $x_0^{d-4}x_1$ but has term x_0^{d-3} and q' does not have monomial x_0^{d-3} but has term $x_0^{d-4}x_1$. This implies that the forms $x_0^2x_2q$ and $x_0^2x_1q'$ vanish at Z.

Let $\mathcal{E}$ be a sheaf over $[\Phi]$. Let C be its support. By Lemma A.2 one concludes that $\mathcal{E}$ is singular at pt_1 if and only if in the homogeneous polynomial of degree d defining C the coefficients in front of the monomials $x_0^{d-2}x_1^2$ and $x_0^{d-1}x_2$ vanish. Therefore, taking into account the considerations above, sheaves over $[\Phi]$ singular at pt_1 constitute a projective subspace of codimension 2 in the fibre $\nu^{-1}([\Phi]) \cong \mathbb{P}_{3d-1}$.

3.2.2. *Sheaves singular at simple points.* Assume in addition that $pt_2 = \langle 0, 0, 1\rangle$. We can still do this without loss of generality because Z can not lie on a line. Then the matrix above is

$$\begin{pmatrix} 1 & 0 & 0 & \dots & 0 \\ 0 & 1 & 0 & \dots & 0 \\ 0 & 0 & 0 & \dots & 1 \\ m_1(pt_3) & m_2(pt_3) & m_3(pt_3) & \dots & m_l(pt_3) \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ m_1(pt_{l-1}) & m_2(pt_{l-1}) & m_3(pt_{l-1}) & \dots & m_l(pt_{l-1}) \end{pmatrix}$$

Then, as in 3.1, we obtain a homogeneous polynomial q'' vanishing at $Z \setminus \{pt_2\}$ with coefficient 1 in front of the monomial $m_l = x_2^{d-3}$. Then the polynomials $x_1x_2^2q''$ and $x_0x_2^2q''$ vanish at Z, the former one has $x_1x_2^{d-1}$ but does not have $x_0x_2^{d-1}$, the latter one has $x_0x_2^{d-1}$ but does not have $x_1x_2^{d-1}$. This means that the sheaves over $[\Phi]$ singular at pt_2 constitute a projective subspace of codimension 2 in the fibre $\nu^{-1}([\Phi]) \cong \mathbb{P}_{3d-1}$. This also shows that the sheaves over $[\Phi]$ singular at pt_j such that pt_1 and pt_j do not lie on a line constitute a projective subspace of codimension 2 in the fibre $\nu^{-1}([\Phi]) \cong \mathbb{P}_{3d-1}$.

Suppose there exists a point pt_j such that pt_1 and pt_j lie on a line. Without loss of generality we can assume that in this case $pt_j = \langle 0, 1, 0\rangle$. Then we can construct a homogeneous polynomial q''' of degree $d-3$ through $Z \setminus \{pt_j\}$ with coefficient 1 in front of the monomial x_1^{d-3}. Then the polynomials $x_0x_1^2q'''$ and $x_1^2x_2q'''$ vanish at Z, the former one has $x_0x_1^{d-1}$ but does not have $x_1^{d-1}x_2$, the latter one has $x_1^{d-1}x_2$ but does not have $x_0x_1^{d-1}$. This means that the sheaves over $[\Phi]$ singular at pt_j such that pt_1 and pt_j lie on a line constitute a projective subspace of codimension 2 in the fibre $\nu^{-1}([\Phi]) \cong \mathbb{P}_{3d-1}$. This concludes the proof of the following.

Lemma 3.4. *The fibres of M_0' over N_1 are unions of $l-1$ different linear subspaces of $\nu^{-1}([\Phi]) \cong \mathbb{P}_{3d-1}$ of codimension 2. In particular the fibres are singular.*

3.3. **Main result.** Now we are able to prove Theorem 1.1.

Singularities. Notice that a generic fibre of a surjective morphism of smooth varieties must be smooth. Therefore, since a generic fibre of M_0' over N is singular as demonstrated in 3.1, we conclude that M_0' is singular. Therefore, M' is a singular subvariety in M.

Dimension. Since, as shown in [10], the codimension of the complement of $\mathbb{B}_0$ in M is at least 2, in order to demonstrate that $\operatorname{codim}_M M' = 2$, it is enough to show that $\operatorname{codim}_{\mathbb{B}_0} \mathbb{B}_0 \cap M' = 2$. Denote $\mathbb{B}_c = \mathbb{B}|_{N_c}$, $\mathbb{B}_1 = \mathbb{B}|_{N_1}$. Since the complement of $N_c \sqcup N_1$ in N_0 has codimension 2, it is enough to show

$$\operatorname{codim}_{\mathbb{B}_c \sqcup \mathbb{B}_1} M' \cap (\mathbb{B}_c \sqcup \mathbb{B}_1) = 2.$$

The latter follows immediately since the codimension of fibres of M'_0 over $N_c \sqcup N_1$ is 2. This concludes the proof.

Smooth locus of M'.

Proposition 3.5. *The smooth locus of M' over N_c coincides with the locus of sheaves corresponding to $Z \subseteq C$ such that only one of the points in Z is a singular point of C.*

Proof. Notice that $H_c \cong N_c$ can be seen as an open subscheme in $S^l\mathbb{P}_2$. Taking the composition of a local section (in analytic or étale topology) of the quotient $\prod_1^l \mathbb{P}_2 \to S^l\mathbb{P}_2$ with the projection $\prod_1^l \mathbb{P}_2 \to \mathbb{P}_2$ to the j-th factor, we get, locally around a given $Z_0 \in H_c$, l different local choices

$$N_c \cong H_c \supseteq U \xrightarrow{p_j} \mathbb{P}_2, \quad j = 1, \dots, l,$$

of one point in $Z \in U \subseteq H_c$. Shrinking U if necessary we can assume that $\mathbb{B} \to N_c$ is trivial over U. Then by 3.1 the subvariety $S_j \subseteq \mathbb{B}|_U$ of those sheaves given by $Z \subseteq C$ that are singular at point $p_j(Z) \in \mathbb{P}_2$ is isomorphic to a product of U with a linear subspace of $\mathbb{P}_{3d-1}$ of codimension 2 (i. e., with $\mathbb{P}_{3d-3}$). Therefore, S_j is smooth. Notice that $M' \cap \mathbb{B}|_U$ is isomorphic to the union of S_j, $j = 1, \dots, l$. Therefore, $\bigcup_j S_j \setminus \bigcup_{j \neq i} S_j \cap S_i$ is smooth, which proves the required statement. □

4. Modifying the boundary by vector bundles

4.1. **Normal spaces at $M' \cap \mathbb{B}_c$.** Let $[\mathcal{F}] \in \mathbb{B}_c \cap M'$ be the isomorphism class of a singular sheaf represented by a curve C of degree d and a configuration of l points $Z \subseteq C$, $Z = \{pt_1, \dots, pt_l\}$. Assume without loss of generality that $Z \cap \operatorname{Sing} C = \{pt_1, \dots, pt_r\} := Z'$, $0 < r \leqslant l$.

Using, for every $j = 1, \dots, r$, the local choice of a point p_j and the morphism $M \to \mathbb{P}S^dV^*$, $[\mathcal{G}] \mapsto \operatorname{Supp} \mathcal{G}$, we obtain locally around $[\mathcal{F}]$ a morphism $\mathbb{B}_c \supseteq U_{[\mathcal{F}]} \xrightarrow{\rho_j} C_d$ from a neigbourhood of $[\mathcal{F}]$ to the universal planar curve C_d of degree d.

This induces a linear map of the tangent spaces $\mathrm{T}_{[\mathcal{F}]} \mathbb{B}_c \to \mathrm{T}_{(C, pt_j)} C_d$. Let C'_d denote the universal singular locus of C_d. Since ρ_j maps S_j to C'_d, we obtain also the induced linear map on the normal spaces

$$N^{(j)}_{[\mathcal{F}]} := \mathrm{T}_{[\mathcal{F}]} \mathbb{B}_c / \mathrm{T}_{[\mathcal{F}]} S_j \xrightarrow{\mathrm{T}_{[\mathcal{F}]}(\rho_j)} \mathrm{T}_{(C, pt_j)} C_d / \mathrm{T}_{(C, pt_j)} C'_d =: N_{(C, pt_j)}.$$

Lemma 4.1. *The linear map $N^{(j)}_{[\mathcal{F}]} \xrightarrow{\mathrm{T}_{[\mathcal{F}]}(\rho_j)} N_{(C, pt_j)}$ constructed above is an isomorphism of 2-dimensional vector spaces.*

Proof. Let $F = \nu^{-1}(\nu([\mathcal{F}]))$ be the fibre of $[\mathcal{F}]$. As already noticed F can be seen as the space of curves of degree d through Z. For a fixed $j \in \{1, \dots, r\}$ let F'_j be the fibre of S_j over $\nu([\mathcal{F}])$, which can be seen as the subspace of F of those curves through Z singular at pt_j. Let F_{pt_j} be the space of curves of degree d through pt_j and let F'_{pt_j} be its subspace of curves singular at pt_j.

Our analysis in 3.1 implies that ρ_j induces an isomorphism $\mathrm{T}_C F / \mathrm{T}_C F'_j \cong \mathrm{T}_C F_{pt_j} / \mathrm{T}_C F'_{pt_j}$, which concludes the proof because both S_j and C'_d are locally trivial over N and $\mathbb{P}_2$ respectively and hence $N^{(j)}_{[\mathcal{F}]} \cong \mathrm{T}_C F / \mathrm{T}_C F'_j$, $N_{(C, pt_j)} \cong \mathrm{T}_C F_{pt_j} / \mathrm{T}_C F'_{pt_j}$. □

4.2. **R-bundles.** Let $\boldsymbol{\mathcal{U}}$ denote the universal family on $M\times\mathbb{P}_2$. Consider a germ of a morphism γ of a smooth curve T to $\mathbb{B}_c$ mapping $0\in T$ to $\gamma(0)=[\mathcal{F}]$. Let $\boldsymbol{\mathcal{F}}$ be the pullback of $\boldsymbol{\mathcal{U}}$ along $\gamma\times\mathrm{id}_{\mathbb{P}_2}$.

If γ is not tangent to S_j at $[\mathcal{F}]$, then $\boldsymbol{\mathcal{F}}$ represents $[\mathcal{F}]$ as a flat 1-parameter degeneration of sheaves $[\boldsymbol{\mathcal{F}}_t]=\gamma(t)$ non-singular at point $p_j(\nu\circ\gamma(t))$, where $\boldsymbol{\mathcal{F}}_t=\boldsymbol{\mathcal{F}}|_{\{t\}\times\mathbb{P}_2}$. If γ is not tangent to S_j at $[\mathcal{F}]$ for all $j=1,\dots,r$, then $\boldsymbol{\mathcal{F}}$ is a degeneration of non-singular sheaves to $\mathcal{F}=\boldsymbol{\mathcal{F}}_0$.

Let $\widetilde{T\times\mathbb{P}_2}\xrightarrow{\sigma}T\times\mathbb{P}_2$ be the blow-up $\mathrm{Bl}_{\{0\}\times Z'}(T\times\mathbb{P}_2)$ and let $D_1=D_1(Z')$ be its exceptional divisor, which is a disjoint union of projective planes $D_1(p_j)\cong\mathbb{P}_2$, $j=1,\dots,r$.

The fibre of the flat morphism $\widetilde{T\times\mathbb{P}_2}\xrightarrow{\sigma}T\times\mathbb{P}_2\xrightarrow{pr_1}T$ over 0 is a reduced surface $D(Z')=D(p_1,\dots,p_r)$ obtained by blowing-up $\mathbb{P}_2$ at $\{pt_1,\dots,pt_r\}$ and attaching the surfaces $D_1(pt_j)\cong\mathbb{P}_2$, $j=1,\dots,r$, to $D_0(Z')=D_0(pt_1,\dots,pt_r)=\mathrm{Bl}_{Z'}\mathbb{P}_2$ along the exceptional lines $L_1,\dots,L_r$.

Let $\boldsymbol{\mathcal{E}}$ be the sheaf on $\widetilde{T\times\mathbb{P}_2}$ obtained as the quotient of the pullback $\sigma^*(\boldsymbol{\mathcal{F}})$ by the subsheaf $\mathit{Tor}_{\mathcal{I}_{D_1}}(\sigma^*(\boldsymbol{\mathcal{F}}))$ generated by the sections annihilated by the ideal sheaf $\mathcal{I}_{D_1}$ of the exceptional divisor D_1.

Lemma 4.2. *Assume that γ is not tangent to S_j at $[\mathcal{F}]$ for all $j=1,\dots,r$. Then the sheaf $\boldsymbol{\mathcal{E}}$ is a flat family of 1-dimensional sheaves. Its fibres $\boldsymbol{\mathcal{E}}_t$, $t\neq 0$, are non-singular sheaves on $\mathbb{P}_2$, the fibre $\mathcal{E}=\boldsymbol{\mathcal{E}}_0$ is a 1-dimensional non-singular sheaf on $D(Z')$.*

Proof. We shall show that our definition of $\boldsymbol{\mathcal{E}}$ locally coincides with the construction from [5].

Fix some local coordinates x_i, y_i at pt_i. Then in some neighbourhood U_i of pt_i the sheaf $\mathcal{F}$ is just an ideal sheaf of pt_i on C and hence can be given as the cokernel of a morphism

$$2\mathcal{O}_{U_i}\xrightarrow{A}2\mathcal{O}_{U_i},\quad A=\left(\begin{smallmatrix}x_i & y_i\\ a_i & b_i\end{smallmatrix}\right),$$

for some polynomials a_i, b_i in x_i, y_i. As $pt_i\in\operatorname{Sing}S$ for $i=1,\dots,r$, a_i and b_i do not have constant terms.

Let $\boldsymbol{\mathcal{U}}_d$ denote the universal family on $C_d\times\mathbb{P}_2$. Then, for a fixed $j\in\{1,\dots,r\}$, locally around the point $[\mathcal{F}]\times pt_j\in\mathbb{B}_c\times\mathbb{P}_2$, $\boldsymbol{\mathcal{U}}$ is isomorphic to the pullback of $\boldsymbol{\mathcal{U}}_d$ along $\rho_j\times\mathrm{id}_{\mathbb{P}_2}$. Then $\boldsymbol{\mathcal{F}}$ is isomorphic locally around $0\times pt_j$ to the pullback of $\boldsymbol{\mathcal{U}}_d$ along $\gamma_j\times\mathrm{id}_{P_2}$, where $\gamma_j=\rho_j\circ\gamma$.

The latter means that the family $\boldsymbol{\mathcal{F}}$ is given around $0\times pt_j$ as a cokernel of the morphism

$$2\mathcal{O}_{T\times U_j}\xrightarrow{\left(\begin{smallmatrix}x_j & y_j\\ v_j & w_j\end{smallmatrix}\right)+t\cdot B(t)}2\mathcal{O}_{T\times U_j},\quad B(t)=\begin{pmatrix}b_{11}(t) & b_{12}(t)\\ b_{21}(t) & b_{22}(t)\end{pmatrix},$$

where $b_{11}(t), b_{12}(t)$ take values in $\Bbbk$ and $b_{21}(t), b_{22}(t)$ are polynomials of degree not bigger than d.

The blow-up $\widetilde{T\times\mathbb{P}_2}$ over $0\times pt_j\in T\times\mathbb{P}_2$ is locally just the blow-up $\mathrm{Bl}_{0\times pt_j}T\times U_j=:\widetilde{T\times U_j}$, it can be seen as a subvariety in $T\times U_j\times\mathbb{P}_2$ given by the (2×2)-minors of the matrix $\left(\begin{smallmatrix}t & x_j & y_j\\ u_0 & u_1 & u_2\end{smallmatrix}\right)$, where u_0, u_1, u_2 are some homogeneous coordinates of the last factor $\mathbb{P}_2$. The canonical section s of $\mathcal{O}_{\widetilde{T\times\mathbb{P}_2}}(D_1)$ is locally given by t/u_0, x_j/u_1, y_j/u_2.

As in [5] one "divides" the pull back $\sigma^*(A+tB(t))$ by s and obtains a family $\boldsymbol{\mathcal{E}}'$ of one-dimensional sheaves given by a locally free resolution

$$0\to 2\mathcal{O}_{\widetilde{T\times U_j}}(D_1)\xrightarrow{\phi(A,B)}2\mathcal{O}_{\widetilde{T\times U_j}}\to\boldsymbol{\mathcal{E}}'\to 0$$

We claim that this construction coincides with taking the quotient by the subsheaf annihilated by $\mathcal{I}_{D_1}$. This follows from a diagram chasing on the following commutative diagram with exact

rows and columns

$$\begin{array}{ccccccccc}
 & & 0 & & & & 0 & & \\
 & & \downarrow & & & & \downarrow & & \\
 & & \mathcal{K} & & & & \mathcal{K} & & \\
 & & \downarrow & & & & \downarrow & & \\
0 & \longrightarrow & 2\mathcal{O}_{\widetilde{T\times U_j}} & \xrightarrow{\sigma^*(A+tB(t))} & 2\mathcal{O}_{\widetilde{T\times U_j}} & \longrightarrow & \sigma^*\boldsymbol{\mathcal{F}} & \longrightarrow & 0 \\
 & & \downarrow{\scriptstyle \left(\begin{smallmatrix} s & 0 \\ 0 & s \end{smallmatrix}\right)} & & \| & & \downarrow & & \\
0 & \longrightarrow & 2\mathcal{O}_{\widetilde{T\times U_j}}(D_1) & \xrightarrow{\phi(A,B)} & 2\mathcal{O}_{\widetilde{T\times U_j}} & \longrightarrow & \boldsymbol{\mathcal{E}}' & \longrightarrow & 0. \\
 & & \downarrow & & & & \downarrow & & \\
 & & \mathcal{K} & & & & 0 & & \\
 & & \downarrow & & & & & & \\
 & & 0 & & & & & &
\end{array}$$

More precisely, one shows that $\mathcal{K}$ is exactly the subsheaf of $\sigma^*\boldsymbol{\mathcal{F}}$ annihilated by s.

Now the flatness of $\boldsymbol{\mathcal{E}}$ and the local freeness of $\mathcal{E}$ on its support follow from [5] (Lemma 5.1 and the discussion before it). □

Remark 4.3. Notice that the fibre of $\sigma^*\boldsymbol{\mathcal{F}}$ over $0 \in T$ is not a 1-dimensional sheaf. Its support contains D_1. The subsheaf $\mathscr{T}or_{\mathcal{I}_{D_1}}(\sigma^*(\boldsymbol{\mathcal{F}}))$ of $\sigma^*(\boldsymbol{\mathcal{F}})$ is the maximal subsheaf supported completely in D_1.

Remark 4.4. The sheaf $\boldsymbol{\mathcal{E}}_0$ depends only on the derivative of γ at 0, i. e., only on the induced map $\mathrm{T}_0\, T \to \mathrm{T}_{[\mathcal{F}]}\, \mathbb{B}_0$ of tangent spaces.

Definition 4.5. *The sheaf $\boldsymbol{\mathcal{E}}_0$ on $D(Z')$ as above is called an R-bundle associated to the pair $Z \subseteq C$, $Z' = Z \cap \operatorname{Sing} C$.*

The following generalizes Definition 5.5 from [5].

Definition 4.6. *Two R-bundles $\mathcal{E}_1$ and $\mathcal{E}_2$ associated to $Z \subseteq C$ on $D(pt_1, \dots, pt_r)$ are called equivalent if there exists an automorphism ϕ of $D(pt_1, \dots, pt_r)$ that acts identically on the surface $D_0(pt_1, \dots, pt_r)$ such that $\phi^*(\mathcal{E}_1) \cong \mathcal{E}_2$.*

Theorem 4.7. *Assume that the components $F_1, \dots, F_r$ intersect transversally at $[\mathcal{F}]$. Then the equivalence classes of R-bundles associated to $Z \subseteq C$ are in one-to-one correspondence with the points of the product of projective lines $\prod_{j=1}^{r} \mathbb{P}N^{(j)}_{[\mathcal{F}]}$.*

Proof. The classes of R-bundles around $D_1(pt_j)$ are parameterized by $\mathbb{P}N_{(C,pt_j)} \cong \mathbb{P}N^{(j)}_{[\mathcal{F}]}$ by [5, Proposition 5.6] and Lemma 4.1. Since the components $F_1, \dots, F_r$ intersect transversally, the map

$$\mathrm{T}_{[\mathcal{F}]}\, \mathbb{B}_c \setminus \cup_{j=1}^{r} \mathrm{T}_{[\mathcal{F}]}\, S_j \to \prod_{j=1}^{r} \mathbb{P}N^{(j)}_{[\mathcal{F}]}$$

is surjective, which means that the class of an R-bundle around a given exceptional plane $D_1(pt_j)$ is independent of the class of the R-bundle around other planes $D_1(pt_i)$, $i \neq j$. This concludes the proof. □

Remark 4.8. The assumption of Theorem 4.7 is always satisfied at least for $r \leqslant 3$ by Remark 3.2. In particular it is the case for $d = 4$. Remark 3.2 also implies that the locus of $[\mathcal{F}] \in \mathbb{B}_c$ that do not satisfy the assumption on transversality lies at least in codimension 7.

Remark 4.9. All possible R-bundles can be produced simultaneously as fibres of a family of sheaves over (an open subset of) $\widetilde{M} := \mathrm{Bl}_{M'} M$. Indeed, pull back the universal family over M to a family $\boldsymbol{\mathcal{U}}_{\widetilde{M}}$ over $\widetilde{M}$. Let $\widetilde{\widetilde{M} \times \mathbb{P}_2} \to \widetilde{M} \times \mathbb{P}_2$ be the blow up along the subvariety in $\widetilde{M} \times \mathbb{P}_2$

where $\boldsymbol{\mathcal{U}}_{\widetilde{M}}$ is singular, pull $\boldsymbol{\mathcal{U}}_{\widetilde{M}}$ back and consider the quotient $\widetilde{\boldsymbol{\mathcal{U}}}$ by the subsheaf annihilated by the ideal sheaf of the exceptional divisor of this blow-up.

Let $\mathbb{B}_{gen}$ denote the open locus of those $[\mathcal{F}] \in \mathbb{B}_0$ satisfying the conditions of Theorem 4.7. Then $\widetilde{\mathbb{B}}_{gen} := \mathrm{Bl}_{M' \cap \mathbb{B}_{gen}} \mathbb{B}_{gen}$ is smooth by Theorem 1.2 and Theorem 1.3 from [7]. The fibre of the exceptional divisor over $[\mathcal{F}] \in \mathbb{B}_{gen}$ coincides with the product $\prod_{j=1}^{r} \mathbb{P}N^{(j)}_{[\mathcal{F}]}$. The restriction of $\widetilde{\boldsymbol{\mathcal{U}}}$ to $\widetilde{\mathbb{B}}_{gen}$ is a flat family of 1-dimensional non-singular sheaves: non-singular $(dm-1)$-sheaves together with R-bundles associated to $Z \subseteq C$, $Z \cap \operatorname{Sing} C \neq \emptyset$.

This allows us to see the blow-up $\widetilde{\mathbb{B}}_{gen}$ as a process that substitutes the boundary $M' \cap \mathbb{B}_{gen}$ by a divisor consisting of non-singular 1-dimensional sheaves, which generalizes the construction from [5].

4.3. **Further work.** A more detailed discussion of the relation of the blow-up $\mathrm{Bl}_{M'} M$ with modifying the boundary of M by vector bundles should follow in a separate paper.

Appendix A. On the ideals of points on planar curves

Let $R = \mathcal{O}_{C,p}$ be a local $\Bbbk$-algebra of a curve C at point $p \in C$. Let $I \subseteq R$ be an ideal of R. As a submodule of a free module, I is a torsion free R-module. If R is regular, i. e., if p is a smooth point of C, then I is free. Therefore, the non-regularity of R is a necessary condition for the non-freeness of I.

A.1. **Ideals of simple points on a curve.** Let $\mathfrak{m} = \mathfrak{m}_{C,p}$ be the maximal ideal of R and let $\Bbbk_p = R/\mathfrak{m}$ be the local ring of the structure sheaf of the one point subscheme $\{p\} \subseteq C$.

Lemma A.1. *Consider an exact sequence of R-modules.*

$$0 \to \mathfrak{m} \to R \to \Bbbk_p \to 0$$

with a non-zero R-module $\mathfrak{m}$. Then $\mathfrak{m}$ is free if and only if R is regular.

Proof. If $\mathfrak{m}$ is free, then $\mathfrak{m} \cong R$ (otherwise $\mathfrak{m} \to R$ would not be injective) and we obtain an exact sequence of R-modules

$$0 \to R \to R \to \Bbbk_p \to 0,$$

which means that the maximal ideal $\mathfrak{m}$ of p is generated by one element. Therefore, R is regular in this case.

Vice versa, assume R is regular. Notice that $\mathfrak{m}$ is always a torsion free R-module as a submodule of R. Therefore, if R is regular, $\mathfrak{m}$ is free as a torsion free module over a regular one-dimensional local ring. □

A.2. **Ideals of fat curvilinear points on a planar curve.** Assume that C is a planar curve locally defined as the zero locus of $f \in \Bbbk[x, y]$. Assume $p = 0$, consider a fat curvilinear point Z at p given by the ideal

$$(x - h(y), y^n) \subseteq \Bbbk[x, y], \quad h(y \in \Bbbk[y]), \quad h(0) = 0, \quad \deg h < n,$$

and assume that Z is a subscheme of C. Let $I \subseteq R$ be its ideal.

Since we assumed $Z \subseteq C$, one can write

$$f = \det \begin{pmatrix} x - h(y) & y^n \\ u(y) & v(x, y) \end{pmatrix}, \quad u(y) \in \Bbbk[y], \quad v(x, y) \in \Bbbk[x, y].$$

Lemma A.2. *Keeping the notations as above, let R be a non-regular ring, i. e., let p be a singular point of C. Then I is non-free if and only if $u(0) = 0$.*

If I is free, then it is generated by $x - h(y)$ and there is an isomorphism

$$R \cong I, \quad r \mapsto r \cdot (x - h(y)).$$

Proof. First of all we shall show that I is generated by one element if and only if $u(0) \neq 0$.

Since $R = \Bbbk[x,y]_{(x,y)}/(f)$, it is enough to answer the question when the ideal

$$(x - h(y), y^n, f) \subseteq \Bbbk[x,y]_{(x,y)}$$

equals (ξ, f) for some $\xi \in \Bbbk[x,y]_{(x,y)}$.

"$\Leftarrow$": If $u(0) \neq 0$, then $(x - h(y), y^n, f) = (x - h(y), f)$.

"$\Rightarrow$": Let $(x - h(y), y^n, f) = (\xi, f)$, without loss of generality we can assume that

$$\xi \in (x - h(y), y^n)$$

and

$$\xi = a \cdot (x - h(y)) + b \cdot y^n$$

for some $a, b \in \Bbbk[x,y]_{(x,y)}$. In order to show that $u(0) \neq 0$, we suppose that the contrary holds true, i. e., $u(0) = 0$.

Since one can embed $\Bbbk[x,y]_{(x,y)}$ into the ring of formal power series $\Bbbk[[x,y]]$, we are going to consider the elements of $\Bbbk[x,y]_{(x,y)}$ as power series in x, y.

Since $x - h(y) \in (\xi, f)$, then $x - h(y) = c \cdot \xi + d \cdot f = ca(x - h(y)) + cby^n + df$. As the orders of cby^n and df are at least 2, we conclude that a and c are units. We can assume without loss of generality that

$$\xi = x - h(y) + \eta(x,y), \quad \operatorname{ord} \eta(h(y), y) \geqslant n.$$

As $y^n \in (\xi, f)$, it must hold $y^n = C\xi + Df$ for some C and D. Since by our assumption $u(0) = 0$, evaluating this equality at $x = h(y)$, we conclude that

$$y^n = C(h(y), y) \cdot \eta(h(y), y) + D(h(y), y) \cdot (-u(y) \cdot y^n).$$

Therefore, C must be a unit and $\operatorname{ord} \xi(h(y), y) = \operatorname{ord} \eta(h(y), y) = n$.

Substituting x by $h(y)$ in the equality $x - h(y) = c \cdot \xi + d \cdot f$, we get

$$0 = c(h(y), y) \cdot \eta(h(y), y) + d(h(y), y) \cdot (-u(y) \cdot y^n).$$

Since c is a unit, it contains a non-zero constant term and hence the product $c(h(y), y) \cdot \eta(h(y), y)$ has order n. On the other hand, since $u(0) = 0$ by our assumption, the order of $d(h(y), y) \cdot u(y) y^n$ is at least $n + 1$. We obtain a contradiction, which shows that our assumption was wrong.

Notice that if I is free, then it must be one-generated. On the other hand we see that if I is one-generated, then it is generated by $x - h(y)$. In this case $u(0) \neq 0$ and therefore f is not divisible by $x - h(y)$. Thus $x - h(y)$ is not a zero divisor in R and there is an isomorphism $R \cong I$, $r \mapsto rx$. The latter means that I is one-generated if and only if it is free, which concludes the proof. □

References

[1] J.-M. Drezet. Fibrés exceptionnels et variétés de modules de faisceaux semi- stables sur $\mathbb{P}_2(\mathbb{C})$. (Exceptional bundles and moduli varieties of semi-stable sheaves on $\mathbb{P}_2(\mathbb{C})$). *J. Reine Angew. Math.*, 380:14–58, 1987.

[2] Jean-Marc Drézet. Variétés de modules alternatives. *Ann. Inst. Fourier (Grenoble)*, 49(1):v–vi, ix, 57–139, 1999.

[3] Geir Ellingsrud and Stein Arild Strømme. On the Chow ring of a geometric quotient. *Ann. of Math. (2)*, 130:159–187, 1989.

[4] Oleksandr Iena. On the singular sheaves in the fine Simpson moduli spaces of 1-dimensional sheaves supported on plane quartics. `ArXiv:1305.2400v2 [math.AG]`, September 2015.

[5] Oleksandr Iena. Universal plane curve and moduli spaces of 1-dimensional coherent sheaves. *Comm. Algebra*, 43(2):812–828, 2015.

[6] J. Le Potier. Faisceaux semi-stables de dimension 1 sur le plan projectif. *Rev. Roumaine Math. Pures Appl.*, 38(7-8):635–678, 1993.

[7] Li Li. Wonderful compactification of an arrangement of subvarieties. *Mich. Math. J.*, 58(2):535–563, 2009.

[8] Mario Maican. On two notions of semistability. *Pac. J. Math.*, 234(1):69–135, 2008.

[9] Carlos T. Simpson. Moduli of representations of the fundamental group of a smooth projective variety. I. *Inst. Hautes Études Sci. Publ. Math.*, (79):47–129, 1994.

[10] Yao Yuan. Moduli spaces of semistable sheaves of dimension 1 on $\mathbb{P}^2$. *Pure Appl. Math. Q.*, 10(4):723–766, 2014.

University of Luxembourg, Campus Kirchberg, Mathematics Research Unit, 6, rue Richard Coudenhove-Kalergi, L-1359 Luxembourg City, Grand Duchy of Luxembourg
E-mail address: oleksandr.iena@uni.lu

University of Luxembourg, Campus Kirchberg, Mathematics Research Unit, 6, rue Richard Coudenhove-Kalergi, L-1359 Luxembourg City, Grand Duchy of Luxembourg
E-mail address: alain.leytem@uni.lu